


Indiana Latinos & Smoking

Smoking is a significant risk factor for cancer, heart disease and stroke, the leading causes of death among Latinos. Lung cancer is the leading cause of cancer deaths among Latinos, with lung cancer deaths about three times higher for Latino men than for Latino women. Stroke and hypertension contribute to cardiovascular disease deaths, as 21% of all coronary heart disease deaths in the U.S. are due to smoking.

Data shown here are adult smoking rates for Latinos in Indiana and some comparisons with the U.S. This data illustrates the great burden smoking places on all Hoosiers.

Smoking Rates among Latinos in Indiana and the U.S.

- The smoking rate for Latinos in Indiana (16.8%) is higher than the U.S. average for Latinos of 14.8%.




Latino Youth Smoking Rates

- Smoking rates among middle school youth do not vary widely between African American students (4.7%) and White students (4.1%), however Latino middle school students smoke at a substantially higher rate of approximately 8.8%.
- Latino high school students have a higher smoking rate than African American high school students – 21.0% of Latino youth smoke while 9.2% African American high school youth report smoking.

Tobacco Prevention and Cessation Commission ~ www.in.gov/isdh/tpc ~ 317.234.1787
www.QuitNowIndiana.com ~ www.indianaquitline.net

Indiana Youth Smoking Rates by Race/Ethnicity, 2010 Indiana Youth Tobacco Survey



Quit Intentions of Hoosier Latinos


- Thirty-two percent (32%) of Latino smokers stated that they plan to stop smoking within next thirty days.
- Nearly 9 out of 10 (89%) current Latino smokers tried to quit in the last twelve months.

Tobacco Prevention and Cessation Commission ~ www.in.gov/isdh/tpc ~ 317.234.1787
www.QuitNowIndiana.com ~ www.indianaquitline.net

Indiana Latinos' Attitudes and Beliefs about smoke-free air policy

- The majority of Latinos in Indiana (92%) believe smoking should not be allowed in the workplace.
- Latinos (70.9%) are more likely to believe that exposure to secondhand smoke is a serious health hazard than Whites (56.2%)
- Latinos are more likely to agree that secondhand smoke is a cause of Sudden Infant Death Syndrome (SIDS) and lung cancer than Whites.

**Percentage of
Smoke-free Homes among Latinos by Smoking Status, 2008
Adult Tobacco Survey**



Sources: 2008 Indiana Adult Tobacco Survey; 2004-2010 Behavioral Risk Factor Surveillance System; HHS, Tobacco Use Among U.S. Racial/Ethnic Minority Groups — African Americans, American Indians and Alaska Natives, Asian Americans and Pacific Islanders, and Latinos: A Report of the Surgeon General, 1998; American Cancer Society, Inc. Cancer Facts and Figures 2010

Tobacco Prevention and Cessation Commission ~ www.in.gov/isdh/tpc ~ 317.234.1787
www.QuitNowIndiana.com ~ www.indianaquitline.net